

Cross-calibration results from observations of the INS **RX J1856**, and the WDs **HZ 43** and **Sirius B**



Vadim Burwitz


Intl. Calibration Working Group
June 14-16 2006, Iceland



Overview

- Absolute Calibration Soft X-rays
 - dependant on model spectra of WDs
 - what models to use?, uncertainties?
- Analyse LETG + HRC-S data HZ 43A, Sirius B and RX J1856.5-3754
 - determine spectral parameters
 - improve effective area.
- Compare obtained photon spectra with EUVE and ROSAT
 - and XMM (work by Frank Haberl)

NLTE pure H model spectrum of HZ43



NLTE pure H model spectrum of HZ43





Fig. 2. NLTE pure hydrogen spectra for $T_{\text{eff}}=50700$ K and $\log g=8.0$, calculated with TMAP using two versions of the bound-free and free-free absorption coefficients, Seaton's approximation (dashed curve) and the full Karszas & Latter (1961) description (solid curve). Both models yield practically identical results at optical/ultraviolet wavelengths, but differ by a factor of 1.33 at 44\AA (vertical dotted line). The insert shows an expanded view. Two further dotted lines mark $\lambda = 1300\text{\AA}$ and $\lambda = 5450\text{\AA}$

RX J1856-37 two temperature bb fit



RX J1856-37 two temperature bb fit





Fig. 4. *Chandra* LETG+HRC spectrum of RX J1856 binned to 0.5\AA , derived with the effective area at wavelengths longwards of the carbon edge corrected to fit an abundance of carbon in the interstellar medium of $a(\text{C}) = 245 - 400$ ppm with 60% in dust grains. The two-temperature fit shown is for 320 ppm, with the upper level of the step indicated for 245 and 400 ppm, too. The fit has a $\chi^2 = 97.7$ for 115 dof (see Sect. 4.5). The model parameters are quoted in Table 2.

Simultaneous fit to RXJ1856 and the WDs



Simultaneous fit to RXJ1856 and the WDs




Fig. 5. Simultaneous fit of RXJ1856, HZ43 A, and Sirius B in the wavelength ranges marked by vertical dotted lines (see Sect. 4.4.2). The LETG spectra binned to 0.5\AA are shown as data points, the corresponding best-fit models as solid curves, and the first-order contributions as dashed curves. The area correction function α is shown at the top. It converts the nominal LETG+HRC-S first-order effective area A^0 of the November 2004 release into the adjusted area A used in this paper. Systematic uncertainties in α are indicated by error bars at 46, 70, 90, and 125\AA . The steps in the count rate spectra of HZ43 A and RXJ1856 at 49 and 69\AA result from the detector gaps. Sirius B was observed off axis and its gaps are located differently (see text).

Parameters obtained from fit


Parameter	Value±Error
(a) HZ43 A ($\lambda = 45 - 160 \text{ \AA}$)	
T_{eff} (K)	51126 ± 660
$\log g$	7.90 ± 0.08
R^2/d^2 (10^{-23})	3.011 ± 0.010
N_{HI} (10^{17} cm^{-2})	8.91 ± 0.37
(b) Sirius B ($\lambda = 74 - 160 \text{ \AA}$)	
T_{eff} (K)	24923 ± 115
$\log g$	$8.6 f^1$
R^2/d^2 (10^{-21})	4.877 ± 0.010
N_{HI} (10^{17} cm^{-2})	6.5 ± 2.0^2
(c) RX J1856 ($\lambda = 15 - 74 \text{ \AA}$)	
kT_{spot} (eV)	62.83 ± 0.41
kT_{star} (eV)	32.26 ± 0.72
R_1/d (km/pc)	0.0378 ± 0.0003
R_2/d (km/pc)	0.1371 ± 0.0010
N_{HI} (10^{20} cm^{-2})	1.10 ± 0.03

Table 2. Parameters of HZ43 A, Sirius B, and RX J1856 based on the simultaneous fit of our model spectra to the LETG+HRC count rate spectra in the wavelength intervals given. The quoted $1-\sigma$ ($\Delta\chi^2 = +1$) errors are correlated and derived from fits with the other parameters for each object kept free. The letter *f* indicates: fixed.



¹ Based on Barstow et al. (2005); Holberg et al. (1998)

² Hébrard et al. (1999). Our fit is required to stay within the $1-\sigma$ error.


Comparison of photon spectra



RX J1856.5-3754: A 'stable' neutron star



Individual fits with
all parameters free:



Simultaneous fits with
 N_H linked:

Model:
tbabs*body

from Frank Haberl

Summary

Context. The absolute calibration of space-borne instruments in the soft X-ray regime rests strongly on model spectra of hot white dwarfs.

Aims. We analyze the *Chandra* LETG+HRC observations of the white dwarfs HZ43 A and Sirius B and of the neutron star RX J185635–3754 with the aim of resolving current uncertainties in the soft X-ray spectral fluxes and photospheric parameters of the three stars. We apply the derived photon spectra to a cross-calibration of the LETG+HRC-S with the short-wavelength *EUVE* spectrometer and the *ROSAT* PSPC.

Methods. We tie HZ43 A to the flux of RXJ1856 in the 44–48Å range and perform a simultaneous least squares fit to the LETG+HRC spectra of the three stars. This allows us to determine an internally consistent set of spectral energy distributions and an empirically derived wavelength-dependent correction to the LETG+HRC-S effective area. We employ NLTE model atmospheres calculated with *TMAP* for the white dwarfs and a two-blackbody model for RXJ1856, tied to the respective optical fluxes.

Results. The two-blackbody model for RXJ1856 features a hot spot on a cooler star and yields $kT_{\text{spot}} = 62.8 \pm 0.4 \text{ eV}$ and $kT_{\text{star}} = 32.3 \pm 0.7 \text{ eV}$ with a stellar radius as seen from infinity of $16.0 \pm 0.1 \text{ km}$ for a distance of 117 pc. For HZ43 A, our fit yields $T_{\text{eff}} = 51126 \pm 660 \text{ K}$ and $\log g = 7.90 \pm 0.080$ (cgs) with anti-correlated errors ($1-\sigma$) which include not only the statistical but also the systematic uncertainties of the fit. HZ43AB displays a previously detected bremsstrahlung component with a temperature $kT \simeq 0.6 \text{ keV}$. For Sirius B, we find $T_{\text{eff}} = 24923 \pm 115 \text{ K}$ for fixed $\log g = 8.6$. The calibration of the short-wavelength *EUVE* spectrometer differs from that of the LETG+HRC-S by $15 \pm 7\%$. The *ROSAT* PSPC is found to be correctly calibrated within a few percent and reports of a major miscalibration are unfounded.

Conclusions. We have obtained improved parameters for RXJ185635–3754, HZ43 A, and Sirius B which fit the observations from the optical to the soft X-ray regime. Our approach allows us to quote their absolute spectral fluxes at selected wavelengths which may aid the calibration of other space-borne instruments.